






Bcl-2 Monoclonal Antibody(6B5)

Cat# A20428PI

INFORMATION


Catalog Number	A20428PI
Size:	50/100 ug
Reactivity	Hu, Ms, Rt, Chk
Application	WB, IF, IHC
Host	Ms
Uniprot	P10415
Immunogen	Synthetic Peptide of Bcl-2
Purity	The antibody was affinity-purified from mouse ascites by affinity-chromatography using specific immunogen.
Formula	PBS, pH 7.4, containing 0.02% sodium azide as Preservative and 50% Glycerol.
Specificity	The antibody detects endogenous Bcl-2 proteins.
Purification	The antibody was affinity-purified from mouse ascites by affinity-chromatography using specific immunogen.
Storage_stability	-20°C/1 year

 <p>Western blot analysis of Hela cells. Molecular weight markers are indicated on the left: 94KD, 66KD, 45KD, 35KD, 26KD, and 20KD. A prominent band is visible at the 26KD position.</p>	<p>Western blot analysis of Hela, diluted at 1:1000</p>
 <p>IHC staining of Human tonsil tissue. The image shows brown staining (likely DAB) and blue counterstain (likely hematoxylin) on a paraffin-embedded tissue section.</p>	<p>IHC staining of Human tonsil tissue paraffin-embedded, diluted at 1:200.</p>
 <p>Bcl-2</p> <p>Western blot analysis showing Bcl-2 protein bands. The image displays four distinct bands of varying intensity.</p>	<p>The picture was kindly provided by our customer.</p>




chicken cell lysis


Western Blot analysis of chicken cell lysis using Antibody diluted at 1:1000




Western blot detection of Bcl-2 in human breast cancer cell line MCF-7(A), MDA-MB-231(B) and Cal51(C) using Bcl-2 mouse mAb (1:2000 diluted). Predicted band size: 26kDa. Observed band size: 26kDa. Picture was kindly provided by our customer from Tianjin Medical University Cancer Institute and Hospital.




Immunohistochemical analysis of paraffin-embedded Human breast cancer. 1, Using Bcl-2 Mouse mAb diluted at 1:200 (4°, overnight). 2, High-pressure and temperature Citric acid, pH6.0 was used for antigen retrieval. 3, Secondary antibody was diluted at 1:200 (room temperature, 50min). Picture was kindly provided by our customer from Tianjin Medical University Cancer Institute and Hospital




Immunohistochemical analysis of paraffin-embedded Human Lymph gland. 1, Antibody was diluted at 1:100(4°,overnight). 2, High-pressure and temperature EDTA, pH8.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 30min).




Immunohistochemical analysis of paraffin-embedded Human Lymph gland. 1, Antibody was diluted at 1:100(4°,overnight). 2, High-pressure and temperature EDTA, pH8.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 30min).




Immunohistochemical analysis of paraffin-embedded Human Fallopian tube. 1, Antibody was diluted at 1:100(4°,overnight). 2, High-pressure and temperature EDTA, pH8.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 30min).



mmunohistochemical analysis of paraffin-embedded Human Fallopian tube. 1, Antibody was diluted at 1:400(4°,overnight). 2, High-pressure and temperature EDTA, pH8.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 30min).



Immunohistochemical analysis of paraffin-embedded Human Fallopian tube. 1, Antibody was diluted at 1:400(4°,overnight). 2, High-pressure and temperature EDTA, pH8.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 30min).



Immunohistochemical analysis of paraffin-embedded Human Fallopian tube. 1, Antibody was diluted at 1:400(4°,overnight). 2, High-pressure and temperature EDTA, pH8.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 30min).