



Rabbit Anti-CD68 Antibody

Cat# A02675BS


Upon receipt, store at -20°C . Avoid repeated freeze.

INFORMATION

Product Name	Anti-CD68 Antibody
Cat. No.	A02675BS
Size	20 µl, 50 µl, 100 µl, 200 µl
Product type	Primary Antibody
Species Reactivity	Ms, Rb
Immunogen Range	21-120/335
Host	Rabbit
Clonality	Polyclonal
Tested applications	Western Blot: 1/500 - 1/2000. ELISA=1:500-1000 IHC-P=1:400-800 IHC-F=1:400-800 Flow-Cyt=1µg/Test ICC=1:100-500 IF=1:200-800
Conjugation	Non-Conjugated
Concentration	1 mg/mL
Purification	Affinity purified by Protein A
Alias	D68; CD68 antigen; CD68 molecule; DKFZp686M18236; GP110; Macrophage antigen CD68 (microsialin); macrosialin; SCARD1; Scavenger receptor class D member 1; macrosialin isoform A precursor; CD68_MOUSE; Macrosialin; Gp110; scavenger receptor class D, member 1; macrophage antigen CD68; LAMP4.
Image	 <p>Sample: Bone (Mouse) Lysate at 40 ug Primary: Anti-CD68 at 1/300 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/10000 dilution</p>



Paraformaldehyde-fixed, paraffin embedded (Rat brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (CD68) Polyclonal Antibody, Unconjugated at 1:400 overnight at 4°C, followed by a conjugated Goat Anti-Rabbit IgG antibody for 90 minutes, and DAPI for nuclei staining.



Tissue/cell: mouse kidney tissue; 4% Paraformaldehyde-fixed and paraffin-embedded;

Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum) at 37°C for 20 min;

Incubation: Anti-CD68 Polyclonal Antibody, Unconjugated 1:200, overnight at 4°C, followed by conjugation to the secondary antibody and DAB staining.

PRODUCT USE LIMITATION

These products are intended for research use only.TRE