



α -tubulin Monoclonal Antibody(8F11)

Cat# A20446PI


Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.

INFORMATION


Product Name	α -tubulin Monoclonal Antibody(8F11)
Cat. No.	A20446PI
Size	50ug, 100ug
Uniprot	Q71U36/P68363
Product type	Primary Antibody
Species Reactivity	Hu, Ms, Rt
Immunogen	Recombinant Protein of Tubulin alpha-1A chain
Target	The antibody detects endogenous α -tubulin protein.
Host	Mouse
Clonality	Monoclonal
Tested applications	WB: 1:5000 IHC: 1:200 IF 1:200
Application	WB, IHC-p, IF, IP
Purification Method	The antibody was affinity-purified from mouse ascites by affinity chromatography using specific immunogen.
Storage buffer	PBS, pH 7.4, containing 0.02% sodium azide as Preservative and 50% Glycerol.
Image	 <p>Western blot analysis of 1) HeLa, 2) Rat Brain Tissue, 3) Mouse Brain Tissue, diluted at 1:5000.</p>



1) Input: Mouse Brain Tissue Lysate 2) IP product: IP dilute 1:200




Immunofluorescence analysis of Human-colon-cancer tissue. 1, α -tubulin Monoclonal Antibody (8F11) (red) was diluted at 1:200(4°C, overnight). 2, Cy3 labeled Secondary antibody was diluted at 1:300(room temperature, 50min). 3, Picture B: DAPI(blue) 10min. Picture A: Target. Picture B: DAPI. Picture C: merge of A+B



Immunofluorescence analysis of Mouse-liver tissue. 1, α -tubulin Monoclonal Antibody (8F11) (red) was diluted at 1:200(4°C, overnight). 2, Cy3 labeled Secondary antibody was diluted at 1:300(room temperature, 50min). 3, Picture B: DAPI(blue) 10min. Picture A: Target. Picture B: DAPI. Picture C: merge of A+B



Immunohistochemical analysis of paraffin-embedded Human-uterus-cancer tissue. 1, α -tubulin Monoclonal Antibody(8F11) was diluted at 1:200(4°C,overnight). 2, Sodium citrate pH 6.0 was used for antibody retrieval(>98°C,20min). 3,Secondary antibody was diluted at 1:200(room temperature, 30min). Negative control was used by secondary antibody only.



Immunohistochemical analysis of paraffin-embedded Rat-kidney tissue. 1, α -tubulin Monoclonal Antibody(8F11) was diluted at 1:200(4°C,overnight). 2, Sodium citrate pH 6.0 was used for antibody retrieval(>98°C,20min). 3,Secondary antibody was diluted at 1:200(room temperature, 30min). Negative control was used by secondary antibody only.

PRODUCT USE LIMITATION

These products are intended for research use only.