Ver.1 Date: 20190312

Technical support: order@acebiolab.com

Phone: 886-3-2870051

ATAT1 Antibody


Cat# A05001PC

Upon receipt, store at -20°C. Avoid repeated freeze.


INFORMATION

Product Name	ATAT1 Antibody
Cat. No.	A05001PC
Size	50 μΙ ,100 μΙ
Product type	Primary Antibody
Species Reactivity	Hu, Ms, Rb
Immunogen	Recombinant Human Alpha-tubulin N-acetyltransferase 1 protein (194-238AA)
Host	Rabbit
Clonality	Polyclonal
Tested applications	WB:1:500-1:5000, IHC:1:200-1:500
Conjugation	Non-Conjugated
Purification Method	>95%, Protein G purified
Concentration	1 mg/mL
Alias	Alpha-tubulin N-acetyltransferase 1 (Alpha-TAT) (Alpha-TAT1) (TAT) (EC 2.3.1.108)
	(Acetyltransferase mec-17 homolog), ATAT1, C6orf134 MEC17
Image	120KD → 90KD → 35KD → 25KD → 20KD → Western Blot Positive WB detected in: Rat kidney tissue, Mouse brain tissue, Mouse liver tissue All lanes: ATAT1 antibody at 6μg/ml Secondary Goat polyclonal to rabbit IgG at 1/50000 dilution





IHC image of ATAT1 Antibody diluted at 1:300 and staining in paraffin-embedded human kidney tissue performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



IHC image of ATAT1 Antibody diluted at 1:300 and staining in paraffin-embedded human liver tissue performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.

=

PRODUCT USE LIMITATION

These products are intended for research use only.

