

# Phospho-IKK $\alpha$ / $\beta$ (Ser176/180) Antibody Sampler Kit

Cat# AK0230

Upon receipt, store at -20°C. Avoid freeze/thaw cycles.

## PRODUCT DESCRIPTION

The NF- $\kappa$ B/Rel transcription factors are present in the cytosol in an inactive state, complexed with the inhibitory I $\kappa$ B proteins. Most agents that activate NF- $\kappa$ B do so through a common pathway based on phosphorylation-induced, proteasome-mediated degradation of I $\kappa$ B. The key regulatory step in this pathway involves activation of a high molecular weight I $\kappa$ B kinase (IKK) complex whose catalysis is generally carried out by three tightly associated IKK subunits. IKK $\alpha$  and IKK $\beta$  serve as the catalytic subunits of the kinase and IKK $\gamma$  serves as the regulatory subunit. Activation of IKK depends upon phosphorylation at Ser177 and Ser181 in the activation loop of IKK $\beta$  (Ser176 and Ser180 in IKK $\alpha$ ), which causes conformational changes, resulting in kinase activation.

## PRODUCT INCLUDES

Cat No.	Product name	Quantity	Applications	Reactivity	Host
A340560	IKK alpha/beta Polyclonal Antibody	20 $\mu$ L	WB, IHC, ELISA	Human, Mouse, Rat	Rabbit
A340241	Phospho-IKK alpha/beta (Ser176/177) Polyclonal Antibody	20 $\mu$ L	WB, IHC, ELISA	Human, Mouse, Rat	Rabbit
A340242	Phospho-IKK alpha/beta (Ser180/181) Polyclonal Antibody	20 $\mu$ L	WB, IHC, ELISA	Human, Mouse, Rat	Rabbit
A1013s	Goat Anti-Rabbit IgG (H+L) (peroxidase/HRP conjugated)	120 $\mu$ L	WB, ELISA	Rabbit	Goat

## PRODUCT USE LIMITATION

These products are intended for research use only.