

# Phospho-MAPK Family Antibody Sampler Kit

Cat# AK0232

Upon receipt, store at -20°C. Avoid freeze/thaw cycles.

## PRODUCT DESCRIPTION

p44/42 MAPK (Erk1/2), SAPK/JNK, and p38 MAPK function in protein kinase cascades that play a critical role in the regulation of cell growth, differentiation, and control of cellular responses to cytokines and stress. p44/42 MAPK is activated by growth and neurotrophic factors. Activation occurs through phosphorylation of threonine and tyrosine residues (Thr202 and Tyr204 in human Erk1) at the sequence T\*EY\* by a single upstream MAP kinase kinase (MEK). SAPK/JNK and p38 MAPK are activated by inflammatory cytokines and by a wide variety of cellular stresses. Activation of SAPK/JNK occurs via phosphorylation at Thr183 and Tyr185 by the dual specificity enzyme SEK/MKK4. Both MKK3 and SEK phosphorylate p38 MAPK on tyrosine and threonine at the sequence T\*GY\* to activate p38 MAP kinase.

## PRODUCT INCLUDES

Cat No.	Product name	Quantity	Applications	Reactivity	Host
<b>A340303</b>	Phospho-P38 (Thr180/Tyr182) Polyclonal Antibody	20µL	WB, IHC, ELISA	Human, Mouse, Rat	Rabbit
<b>A340370</b>	Phospho-ERK 1/2 (Thr202/Tyr204) Polyclonal Antibody	20µL	WB, IHC, ELISA	Human, Mouse, Rat	Rabbit
<b>A340251</b>	Phospho-JNK1/2/3 (Thr183/Y185) Polyclonal Antibody	20µL	WB, ELISA	Human, Mouse, Rat	Rabbit
<b>A1013s</b>	Goat Anti-Rabbit IgG (H+L) (peroxidase/HRP conjugated)	120µL	WB, ELISA	Rabbit	Goat

## PRODUCT USE LIMITATION

These products are intended for research use only.