

Smad2/3 Antibody Sampler Kit

Cat# AK0265

Upon receipt, store at -20°C. Avoid freeze/thaw cycles.

PRODUCT DESCRIPTION

Transforming growth factor- β (TGF- β) superfamily signaling plays a critical role in the regulation of cell growth, differentiation, and development in a wide range of biological systems. In general, signaling is initiated with ligand-induced oligomerization of serine/ threonine receptor kinases and phosphorylation of the cytoplasmic signaling molecules Smad2 and Smad3 for the TGF- β /activin pathway, or Smad1/5/8 for the bone morphogenetic protein (BMP) pathway. Carboxy-terminal phosphorylation of Smad proteins by activated receptors results in their partnering with the common signaling transducer Smad4, and translocation to the nucleus. Activated Smad proteins regulate diverse biological effects by partnering with transcription factors resulting in cell-state specific modulation of transcription.

PRODUCT INCLUDES

Cat No.	Product name	Quantity	Applications	Reactivity	Host
A340674	SMAD2 Polyclonal Antibody	20 μ L	WB, IHC, ELISA	Human, Mouse, Rat	Rabbit
A340328	Phospho-SMAD2 (Ser467) Polyclonal Antibody	20 μ L	WB, IHC, ELISA	Human, Mouse, Rat	Rabbit
A340675	SMAD2/3 Polyclonal Antibody	20 μ L	WB, IHC, ELISA	Human, Mouse, Rat, Monkey	Rabbit
A340306	Phospho-SMAD2/3 (Thr8) Polyclonal Antibody	20 μ L	WB, ELISA	Human, Mouse, Rat	Rabbit
A340726	SMAD3 Polyclonal Antibody	20 μ L	WB, IHC-p	Human, Mouse, Rat	Rabbit
A340329	Phospho-SMAD3 (Ser425) Polyclonal Antibody	20 μ L	WB, IHC, ELISA	Human, Mouse, Rat	Rabbit
A340734	SMAD4 Polyclonal Antibody	20 μ L	WB, IHC-p, IF, ELISA	Human, Mouse, Rat, Monkey	Rabbit
A1013s	Goat Anti-Rabbit IgG (H+L) (peroxidase/HRP conjugated)	120 μ L	WB, ELISA	Rabbit	Goat

PRODUCT USE LIMITATION

These products are intended for research use only.