





PR Monoclonal Antibody(Z15)



Cat# A630003

Upon receipt, store at -20°C. Avoid repeated freeze.

INFORMATION

Product Name	PR Monoclonal Antibody(Z15)	
Cat. No.	A630003	
Size	100µg/50µg	
Uniprot	Human P06401	
Product type	Primary antibody	
Species Reactivity	Human,Mouse,Rat	
Immunogen	Synthetic Peptide of PR	
Host	Mouse	
Clonality	Monoclonal	
Tested applications	IHC-p,IF	
Application	IHC: 1:200 IF: 1:100-200	
Purification Method	The antibody was affinity-purified from mouse ascites by affinity-chromatography using epitope-specific immunogen.	
Storage instruction	-20°C/1 year	
Alias	PGR; NR3C3; Progesterone receptor; PR; Nuclear receptor subfamily 3 group C member 3	
Image		<p>Immunohistochemical analysis of paraffin-embedded Human-uterus tissue.</p> <ol style="list-style-type: none"> 1,PR Monoclonal Antibody(Z15) was diluted at 1:200(4°C,overnight). 2, Sodium citrate pH 6.0 was used for antibody retrieval(>98°C,20min). 3,Secondary antibody was diluted at 1:200(room temperature, 30min). <p>Negative control was used by secondary antibody only.</p>

		<p>Immunohistochemical analysis of paraffin-embedded Rat-testis tissue.</p> <ol style="list-style-type: none"> 1, PR Monoclonal Antibody(Z15) was diluted at 1:200(4°C, overnight). 2, Sodium citrate pH 6.0 was used for antibody retrieval(>98°C, 20min). 3, Secondary antibody was diluted at 1:200(room temperature, 30min). <p>Negative control was used by secondary antibody only.</p>
		<p>Immunohistochemical analysis of paraffin-embedded Mouse-testis tissue.</p> <ol style="list-style-type: none"> 1, PR Monoclonal Antibody(Z15) was diluted at 1:200(4°C, overnight). 2, Sodium citrate pH 6.0 was used for antibody retrieval(>98°C, 20min). 3, Secondary antibody was diluted at 1:200(room temperature, 30min). <p>Negative control was used by secondary antibody only.</p>
		<p>Immunofluorescence analysis of Human-appendix tissue.</p> <ol style="list-style-type: none"> 1, PR Monoclonal Antibody(Z15)(red) was diluted at 1:200(4°C, overnight). 2, Cy3 labeled Secondary antibody was diluted at 1:300(room temperature, 50min). 3, Picture B: DAPI(blue) 10min. <p>Picture A: Target. Picture B: DAPI. Picture C: merge of A+B</p>
		<p>Immunofluorescence analysis of Mouse-liver tissue.</p> <ol style="list-style-type: none"> 1, PR Monoclonal Antibody(Z15)(red) was diluted at 1:200(4°C, overnight). 2, Cy3 labeled Secondary antibody was diluted at 1:300(room temperature, 50min).

		<p>3, Picture B: DAPI(blue) 10min. Picture A:Target. Picture B: DAPI. Picture C: merge of A+B</p>
		<p>Immunofluorescence analysis of Rat-heart tissue. 1,PR Monoclonal Antibody(Z15)(red) was diluted at 1:200(4°C,overnight). 2, Cy3 labeled Secondary antibody was diluted at 1:300(room temperature, 50min). 3, Picture B: DAPI(blue) 10min. Picture A:Target. Picture B: DAPI. Picture C: merge of A+B</p>
		<p>Immunohistochemical analysis of paraffin-embedded human breast carcinoma using PR Monoclonal Antibody.</p>

PRODUCT USE LIMITATION

These products are intended for research use only.