






# CRP Polyclonal Antibody



Cat# A22023PI

Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.

## INFORMATION

<b>Product Name</b>	CRP Polyclonal Antibody	
<b>Cat. No.</b>	A22023PI	
<b>Size</b>	20 ug / 50 ug / 100 ug	
<b>Uniprot</b>	P02741	
<b>Product type</b>	Primary antibody	
<b>Species Reactivity</b>	Hu, Rt	
<b>Immunogen</b>	The antiserum was produced against synthesized peptide derived from the Internal region of human CRP. AA range:101-150	
<b>Target</b>	The antibody detects endogenous CRP protein	
<b>Host</b>	Rabbit	
<b>Concentration</b>	1 mg/ml	
<b>Clonality</b>	Monoclonal	
<b>Tested applications</b>	WB: 1:500 – 1:2000; IHC: 1:100 – 1:300; ELISA: 1:20000	
<b>Application</b>	WB, IHC-p, ELISA	
<b>Conjugation</b>	Non-conjugated	
<b>Purification Method</b>	The antibody was affinity-purified from mouse ascites by affinity-chromatography using specific immunogen.	
<b>Storage buffer</b>	PBS, pH 7.4, containing 0.02% sodium azide as Preservative and 50% Glycerol.	
<b>Alias</b>	CRP; PTX1; C-reactive protein	
<b>Image</b>		Western Blot analysis of mouse liver cells using CRP Polyclonal Antibody. Secondary antibody was diluted at 1:20000

		<p>Immunohistochemical analysis of paraffin-embedded human-liver, antibody was diluted at 1:100</p>
		<p>Immunohistochemical analysis of paraffin-embedded rat-liver, antibody was diluted at 1:100</p>
		<p>Immunohistochemical analysis of paraffin-embedded rat-liver, antibody was diluted at 1:100</p>
		<p>Immunohistochemical analysis of paraffin-embedded Human Liver. 1, Antibody was diluted at 1:200(4°, overnight). 2, High-pressure and temperature EDTA, pH8.0 was used for antigen retrieval. 3, Secondary antibody was diluted at 1:200(room temperature, 30min).</p>

		<p>Immunohistochemical analysis of paraffin-embedded Human Liver. 1, Antibody was diluted at 1:200(4°, overnight). 2, High-pressure and temperature EDTA, pH8.0 was used for antigen retrieval. 3, Secondary antibody was diluted at 1:200(room temperature, 30min).</p>
		<p>Immunohistochemical analysis of paraffin-embedded Human Liver. 1, Antibody was diluted at 1:200(4°, overnight). 2, High-pressure and temperature EDTA, pH8.0 was used for antigen retrieval. 3, Secondary antibody was diluted at 1:200(room temperature, 30min).</p>

### **PRODUCT USE LIMITATION**

These products are intended for research use only.